

Equivariant Singular Riemann-Roch Theorem

Bin Zhang

February 1, 2008

1 Introduction

The Riemann-Roch theorem [7], [2] has played such a key role in the study of algebraic geometry, algebraic topology or even number theory, that many attempts have been made to generalize it to the equivariant case (for example [9], [4]). Here we consider the case of the equivariant singular Riemann-Roch theorem, following the framework of P. Baum, W. Fulton and R. MacPherson.

2 Totaro's approximation of EG

The purpose of this section is to describe a construction, due to Totaro [5], of an algebro-geometric substitute for the classifying space of topological group.

For topological group G , the spaces EG and BG are infinite dimensional, it is hard to control in general. What Totaro produces for a reductive algebraic group is a directed system of G -bundles $E_n \rightarrow B_n$ with the following property: for any algebraic principal G -bundle $E \rightarrow X$, there is a map $X' \rightarrow X$ with the fiber isomorphic to affine space \mathbb{A}^m , such that the pullback bundle $E' \rightarrow X'$ is pulled back from one of the bundles in the directed system by a map $X' \rightarrow B_n$.

For a complex linear reductive algebraic group G , Totaro constructs the directed system as follows:

Every object in the directed system is a pair (V, V') , such that V is a representation of G , V' is a non-empty open set of V , G acts freely on V' , $V' \rightarrow V'/G$ is a principal G -bundle. $(V, V') < (W, W')$ if V is a subspace of W , $V' \subset W'$, $\text{codim}_V(V - V') < \text{codim}_W(W - W')$. The morphisms are just inclusions.

For some groups, there is a convenient choice of the directed system. For example, if $G = T$ is a split torus of rank n , then $\{(V^n, (V - \{0\})^n) \mid \dim V = l\}$ is the directed system.

3 Equivariant cohomology

In general [1], we define

$$H_G^*(X) = H^*(X \times^G EG),$$

Using the directed system above,

LEMMA 3.1 *For a complex Lie group G and for X a complex algebraic variety on which G operates algebraically, we have*

$$H_G^*(X) = \varinjlim_{(V, V')} H^*(X \times^G V')$$

Proof. See [3] ■

For every G -vector bundle E over X , $E \times^G V' \rightarrow X \times^G V'$ and $E \times^G EG \rightarrow X \times^G EG$ are all G -vector bundles, so we can define the Chern character $ch^G(E) \in H_G^*(X)$ and $ch^V(E) \in H^*(X \times^G V')$ by

$$ch^G(E) = ch(E \times^G EG)$$

and

$$ch^V(E) = ch(E \times^G V')$$

We know that under the map $H_G^*(X) \rightarrow H^*(X \times^G V')$, $ch^G(E)$ goes to $ch^V(E)$.

When X is smooth, then for any (V, V') in the directed system, $X \times^G V'$ is smooth. So we can define the cohomological equivariant Todd class $Td^G(X)$ to be the element in $H_G^*(X)$ which maps to $Td(X \times^G V')$ for every (V, V') in the directed system.

4 Equivariant homology

Now for a complex linear reductive algebraic group G , by using the constructed directed system, we can define the equivariant homology [5] for a G -variety X , suppose $\dim_{\mathbb{C}} X = n$.

$$H_i^G(X) = \lim_{\leftarrow (V, V')} H_{i+2l-2g}^{BM}(X \times^G V')$$

where $l = \dim_{\mathbb{C}} V$, $g = \dim_{\mathbb{C}} G$.

Remark It is possible that $H_i^G(X) \neq 0$ for negative i .

PROPOSITION 4.1 $H_*^G(X)$ is independent of (V, V') if $\text{codim}_V(V - V')$ big enough.

$H_*^G(X)$ is a module over $H_G^*(X)$. In other words there is a cap product: $\cap : H_G^*(X) \otimes H_*^G(X) \rightarrow H_*^G(X)$ defined in the following way:

If $\alpha \in H_G^*(X)$, then for any (V, V') , the image of α under the map $H_G^*(X) \rightarrow H^*(X \times^G V')$ defines a map by cap product: $H_*^{BM}(X \times^G V') \rightarrow H_*^{BM}(X \times^G V') \rightarrow H_*^G(X)$. Go over to the direct limit, these maps induce a map from $H_*^G(X)$ to $H_*^G(X)$. This gives us the module structure.

For a non-singular variety X , we can define the orientation class $[X]_G \in H_{2n}^G(X)$ as follows:

For any object (V, V') in the directed system, $X \times^G V'$ is non-singular, so we have a class $[X \times^G V'] \in H_{2n+2l-2g}^{BM}(X \times^G V')$, $\{[X \times^G V']\}$ gives the orientation class $[X]_G$.

5 Equivariant Riemann-Roch Theorem

Now we can define the equivariant Todd class τ^G as follows,

For any equivariant coherent G -sheaf \mathcal{F} over X (X is a complex algebraic variety), if $p : X \times V' \rightarrow X$ is the projection, then the pullback sheaf $p^*\mathcal{F}$ is a sheaf over $X \times V'$, it descends to a coherent sheaf \mathcal{F}_V over $X \times^G V'$. $\tau(\mathcal{F}_V)$ is in $H_*^{BM}(X \times^G V')$, so we get a map $K_0^G(X) \rightarrow H_*^{BM}(X \times^G V')$. Obviously, it is compatible with the transition maps, so we get a map $K_0^G(X) \rightarrow H_*^G(X)$, that is τ^G .

THEOREM 5.1 (Equivariant Riemann-Roch Theorem) τ^G is the unique natural transformation (with respect to G -equivariant proper maps) from $K_0^G(X)$ to $H_*^G(X)$, such that for any G -variety X , we have the following commutative diagram:

$$\begin{array}{ccccc} K_G^0(X) & \otimes & K_0^G(X) & \xrightarrow{\otimes} & K_0^G(X) \\ & & \downarrow ch^G \otimes \tau^G & & \downarrow \tau^G \\ H_G^*(X) & \otimes & H_*^G(X) & \xrightarrow{\cap} & H_*^G(X) \end{array}$$

and if $f : X \rightarrow Y$ is a proper G -equivariant algebraic map, then

$$\begin{array}{ccc} K_0^G(X) & \xrightarrow{f_*} & K_0^G(Y) \\ \downarrow \tau^G & & \downarrow \tau^G \\ H_*^G(X) & \xrightarrow{f_*} & H_*^G(Y) \end{array}$$

Furthermore, if X is non-singular, and \mathcal{O}_X is the structure sheaf, then

$$\tau^G(\mathcal{O}_X) = Td^G(X) \cap [X]_G$$

Proof. In fact, the proof is already encoded in the definition. ■

From the discussion, we have the following commutative diagram:

$$\begin{array}{ccc} K_0^G(X) & \xrightarrow{\tau^G} & H_*^G(X) \\ \downarrow & & \downarrow \\ K_0(X) & \xrightarrow{\tau} & H_*(X) \end{array}$$

where both downarrows are induced by forgetting functors.

6 Equivariant Riemann-Roch theorem with value in equivariant Chow group

For linear algebraic group G , by using the same directed system, we can define [6]

$$A_i^G(X) = \lim_{\rightarrow} A_{i+2l-2g}(X \times^G V')$$

where $l = \dim_{\mathbb{C}} V$, $g = \dim_{\mathbb{C}} G$.

For the equivariant operational Chow group $A_G^*(X)$, we define it as

$$A_G^i(X) = \lim_{\leftarrow} A^i(X \times^G V')$$

This definition concises with the definition in [6].

As in the previous section, we can define the Chern character, the Todd class and get a theorem similar to Theorem 5.1, except we use $A_G^i(X) \otimes \mathbb{Q}$ and $A_*^G(X) \otimes \mathbb{Q}$ to replace $H_*^G(X)$ and $H_*^G(X)$.

References

- [1] M.F. Atiyah & R. Bott, The moment map and the equivariant cohomology, *Topology*, Vol. 23, No. 1, 1984, 1-28.
- [2] P. Baum, W. Fulton & R. MacPherson, Riemann-Roch for singular varieties, *Publ. Math. IHES (Paris)* 45 (1975), 101-167.
- [3] M. Brion, Equivariant cohomology and equivariant intersection theory. Notes by Alvaro Rittatore. NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Representation theories and algebraic geometry (Montreal, PQ, 1997), 1-37, Kluwer Acad. Publ., Dordrecht, 1998
- [4] M. Brion & M. Vergne, An equivariant Riemann-Roch theorem for complete, simplicial toric varieties. *J. Reine Angew. Math.* 482 (1997), 67-92.
- [5] D. Edidin & W. Graham, Characteristic classes in the Chow ring, *J. Algebraic Geometry* 6 (1997), 431-443
- [6] D. Edidin & W. Graham, Equivariant intersection theory, preprint
- [7] F. Hirzebruch, Topological methods in algebraic geometry, *Die Grundlehren der Mathematischen Wissenschaften, Band 131* Springer-Verlag New York, Inc., New York 1966
- [8] G. B. Segal, Equivariant K-Theory, *Publ. Math. IHES (Paris)* 34 (1968), 129-151.
- [9] T. Kawasaki, The Riemann-Roch theorem for complex V -manifolds. *Osaka J. Math.* 16 (1979), no. 1, 151-159